

A Comparison between Primary and Secondary Breast Angiosarcoma: Our Local Experience

Chan Tin Ho T, Law Hei Yu H, Au-Yeung Chung Ling K, Ling Y, Lee Suet Ying J



WJMER

World Journal of Medical Education and Research

An Official Publication of the Education and Research Division of Doctors Academy



The Impact of the Covid-19 Pandemic on Undergraduate Medical Education in Namibia: The Educators' Perspective

Case Presentation—the Key Which Unlocks the Diagnosis

A Rare Case of Closed-Loop Small Bowel Obstruction Secondary to Herniation Through A Defect in the Broad Ligament of the Uterus- A Case Report

Survey of Medical Students' Opinions Regarding their Training in the Discipline of Infectious Diseases

A Comparison between Primary and Secondary Breast Angiosarcoma: Our Local Experience

Evaluating Medical Students', Clinicians', and the Public's Perspectives on the Integration of Medical Associate Professionals (MAPs) into the NHS

The Hidden Dangers of Fast-Food and Processed Treats: What Everyone Needs to Know



ISSN 2052-1715



A Comparison between Primary and Secondary Breast Angiosarcoma: Our Local Experience

Chan Tin Ho T¹, Law Hei Yu H², Au-Yeung Chung Ling K¹, Ling Y¹, Lee Suet Ying J²

Institution

¹Department of Surgery,
Ruttonjee Hospital, Hong
Kong

²Department of Surgery,
Pamela Youde Nethersole
Eastern Hospital, Hong
Kong

Abstract:

Breast angiosarcoma is a rare form of cancer and can be classified into primary or secondary types. We report two cases of breast angiosarcoma in our locality, and illustrate the differences encountered between the two types. There is no standard treatment for this disease. Surgery is the mainstay of treatment, but the role of adjuvant treatment remains uncertain.

Key Words:

Breast Angiosarcoma, Surgery, Case Report

Corresponding Author:

Dr Timothy Chan Tin Ho; E-mail: acl768@ha.org.hk

**WJMER, Vol 31: Issue 1,
2025**

Introduction

Angiosarcoma of the breast is a rare and highly aggressive cancer, accounting for only 0.04% of all breast malignancy¹. It can occur as a primary lesion of the breast or as a secondary lesion after breast irradiation or in patients with chronic lymphedema after axillary dissection (Stewart - Treves syndrome). Diagnosis is difficult and the prognosis is poor, with optimal adjuvant treatment based mainly on expert opinion due to the rarity of the condition². We report two cases in our locality and compare the differences encountered between primary and secondary disease.

Case I

Miss C is a 46-year-old lady, whose mother was diagnosed with breast cancer at age of 56, presented with progressive right breast engorgement and mastalgia in 2021. Physical examination revealed breast asymmetry with an 8cm firm mass over the upper outer quadrant of the right breast which was not fixed to the underlying muscle or overlying skin. The nipple areolar complex was not affected and there were no palpable axillary lymph nodes. Ultrasound and bilateral mammogram (figure 1) was performed which showed a large 7.9x 3.1 x 8.8cm irregular mass at the right upper outer quadrant with no suspicious microcalcification and no axillary lymphadenopathy bilaterally. Core biopsy of the lesion showed atypical vascular proliferation, most in keeping with well differentiated angiosarcoma. A whole-body positron emission tomography computed tomography (PET-CT) showed no distant metastasis. After discussion with

the multi-disciplinary team, Miss C underwent a right skin sparing mastectomy with transverse rectus abdominis myocutaneous (TRAM) flap reconstruction and sentinel lymph node biopsy. Pathology confirmed angiosarcoma with clear margins (most margins >2cm), and sentinel lymph node was negative for malignancy. Adjuvant radiotherapy was offered for improving local control based on retrospective series, but Miss C was not keen.

In December 2022, she presented with haemoptysis with ~5mL fresh blood with clots during each coughing episode. On examination, a discrete 2cm right upper jugular cervical lymph node was palpable, and a plain chest radiograph revealed a 18mm opacity over the left hilar region. Ultrasound-guided fine needle aspiration of the enlarged cervical lymph node showed atypical cells. Excisional biopsy of the lymph node was performed and the resulting pathological findings were consistent with metastatic angiosarcoma. Her PET-CT revealed multiple new hypermetabolic bilateral lung nodules with adjacent ground glass densities which were suggestive of haemorrhagic metastasis of angiosarcoma. Within a month of her recurrence, she also developed multiple scalp, gum and skin nodules (figure 2) with occasional bleeding. She was started on palliative chemotherapy, initially with paclitaxel which was later changed to adriamycin and zometa after she developed bone and liver metastasis. She then developed brain metastasis for which she was given radiotherapy (figure 3).

Case 2

In comparison, Madam S is a 76-year old lady with a right breast ductal carcinoma in situ in 2010, for which she underwent a right sided breast-conserving therapy, sentinel lymph node biopsy with adjuvant radiotherapy. In 2014, she had a screen-detected lesion for which she underwent left sided breast-conserving therapy and sentinel lymph node biopsy (pT1cN0 ER/PR+ invasive ductal carcinoma, margins clear) with subsequent radiotherapy and hormonal treatment. Post treatment she has been kept on regular surveillance with no recurrence seen on her latest mammogram and ultrasound in January 2021.

Other significant medical history include papillary thyroid cancer, for which she received a total thyroidectomy and adjuvant radioactive iodine therapy.

In November 2022, Madam S presented with a left breast nodule with associated pain and bleeding. Clinically there was a 1x1cm round nodule at the left 5 o'clock region, 2cm from the nipple with easy contact bleeding. Physical examination revealed no other suspicious breast lump or palpable axillary lymphadenopathy. There was also no lymphoedema. The working diagnosis at the time was a bleeding pyogenic granuloma. In view of the persistent troublesome bleeding and history of bilateral breast cancer, the lesion was excised under local anaesthetic. The pathology was consistent with moderately differentiated angiosarcoma with the close margins (closest 1mm). A PET-CT was performed which showed only the hypermetabolic skin lesion over the left breast related to excisional biopsy (figure 4), with no nodal or distant metastasis. Madam S later underwent wide local excision with no axillary treatment in private in January 2023. The pathology showed no residual tumour. No adjuvant therapy for her breast angiosarcoma was required, and there was no evidence of recurrence at her latest follow up with oncology.

Discussion

Angiosarcoma of the breast is extremely rare and can be divided into the de-novo (primary) or therapy-related (secondary) form. Primary breast angiosarcoma usually occurs between females between 20-50 years of age^{3,4,5}, whilst secondary breast angiosarcoma is typically seen in older women, with the mean age of 70⁶. Secondary cases occur following a latent period after radiation or in patients with chronic lymphedema after axillary dissection (Stewart - Treves syndrome). In contrast, there are no known risk factors for the primary form. High grade breast angiosarcomas are associated with a poor prognosis, tend to

metastasize early, often to the lungs or liver and have a 5-year survival rate of only 15%⁷. Low grade and intermediate grade tumours fare better in some studies⁷, however according to Nasciemento et al, there is no correlation between histologic grade and patient outcome in their review of 49 cases⁸.

Patients with breast angiosarcoma can present with diffuse breast enlargement or a rapidly growing palpable breast mass which may have a tinge of purple discolouration, due to its highly vascular nature. Diagnosing breast angiosarcoma can be difficult as the radiological findings are non-specific. Tumours may appear hypoechoic, hyperechoic or heterogeneous on ultrasound, with high vascularity⁹. Mammographically, breast angiosarcomas can appear as large dense homogenous mass with no calcifications or spiculations⁹ and up to 33% of the tumours can have normal mammogram findings¹⁰. Magnetic resonance imaging (MRI) or computed tomography (CT) with contrast can help confirm the hypervascular nature of these lesions. Primary angiosarcomas are often located deep in the breast parenchyma¹¹ hence pathological confirmation with core biopsy may have false negative rates of up to 40% of cases¹².

Currently there is no standard treatment for breast angiosarcoma in view of the small number of reported cases. Surgical resection with negative margins is typically performed, with preference for mastectomy over wide local excision due to lower estimated recurrence rate (8% vs 23% respectively)¹³. Axillary dissection is usually not required as lymph node involvement is uncommon (<5% of cases)¹⁴. Ragavan et al 2020 performed a retrospective review between 2006 and 2019 on axillary lymph node dissection in breast angiosarcoma patients in Singapore with no positive lymph node metastasis detected in their cohort of thirteen cases¹⁵. The role of chemotherapy and radiotherapy in breast angiosarcomas is not well established. For primary breast angiosarcoma, adjuvant radiotherapy after surgery allows for better local control with lower recurrence rate according to Johnstone et al.¹⁶ but the overall survival rate was comparable with those without radiotherapy¹⁷. The role of adjuvant chemotherapy is undefined at present, being based mainly on adult soft tissue sarcomas. Current literature suggests that patients with large high grade angiosarcoma or recurrence may benefit from chemotherapy¹⁸ as these subset of patients have a poor prognosis. Anti-angiogenic and immunological therapies have also been described, but to date have had disappointing results¹⁹.

Conclusion

In conclusion, we report two cases of breast angiosarcoma in our locality. Given the aggressive

nature of the disease, early detection and prompt surgical treatment are needed. Further studies are needed to guide adjuvant treatment modalities and their survival benefit. In addition, it is still unclear whether treatment for primary vs secondary angiosarcomas differ given the rarity of the disease.

Reference

1. Agarwal PK, Mehrotra R. Haemangiosarcoma of the breast, *Ind J Cancer*, 1977; 14(2):182-185.
2. Kim YJ, Ryu JM, Lee SK et al. Primary angiosarcoma of the Breast: A single-centre retrospective study in Korea. *Curr Oncol*. 2022, 29(5), 3272-3281.
3. Sher T, Hennessy BT, Valero V et al. Primary angiosarcoma of the breast. *Cancer* 2007, 110 (1):173-178.
4. Wang XY, Jakowski J, Tawfik OW, Thomas PA, Fan F. Angiosarcoma of the breast: a clinicopathologic analysis of cases from the last 10 years. *Ann Diag Path* 2009. 13(3):147-150.
5. Scow JS, Reynolds CA, Degnim AC, et al. Primary and secondary angiosarcoma of the breast: the Mayo Clinic experience. *J Surg Oncol* 2010. 101(5):401-407.
6. orburger SA, Xing Y, Hunt KK, et al. Angiosarcoma of the breast. *Cancer* 2005. 104 (12):2682-8.
7. Rosen PP, Kimmel M, Ernsberger D. Mammary angiosarcoma. The prognostic significance of tumour differentiation. *Cancer* 1988. 62 (10):2145-51.
8. Nascimento AF, Raut CP, Fletcher CD. Primary angiosarcoma of the breast: clinicopathologic analysis of 49 cases, suggesting that grade is not prognostic. *Am J Surg Pathol*. 2008;32(12):1896-904.
9. Champeaux-Orange E, Bonneau C, Raharimanana B et al. Primary breast angiosarcoma: two case reports. *Cancer Radiother*. 2009;13(3):209-12.
10. Liberman L, Dershaw DD, Kaufman RJ, Rosen PP. Angiosarcoma the breast. *Rad*. 1992. 183(3):649-54.
11. Fletcher CDM, MacGrogan G, Fox SB. Angiosarcoma. In Lakhani SR, Ellis IO, Schnitt SJ, Tan PH, van de Vijver MJ, editors. *World Health Organization classification of tumours of the breast*. Lyon: IARC; 2012. p.135-6
12. Chen KTK, Kirkegaard DD, Bocian JJ. Angiosarcoma of the breast. *Cancer*. 1980;46:268-271.
13. Johnson CM, Garguilo GA. Angiosarcoma of the breast: A case report and literature review. *Curr Surg*. 2002 Sep-Oct;59(5):490-4.
14. Guerrero MA, Ballard BR, Grau AM. Malignant phyllodes tumour of the breast: review of the literature and case report of stromal overgrowth. *Surg Oncol*. 2003;12(1):27-37
15. Ragavan S, Lim HJ, Tan JWS et al. Axillary lymph node dissection in angiosarcomas of the breast: an Asian institutional perspective. *Sarcoma*. 2020: 4890803
16. Johnstone PA, Pierce LJ, Merino MJ, Yang JC, Epstein AH, DeLaney TF. Primary soft tissue sarcomas of the breast: local-regional control with post-operative radiotherapy. *Int J Radiat Oncol Biol Phys*. 1993;27:671-5.
17. Pandey M, Mathew A, Abraham EK, Rajan B. Primary sarcoma of the breast. *J Surg Oncol*. 2004;87:121-5.
18. Sher T, Hennessy BT, Valero V, et al. Primary angiosarcomas of the breast. *Cancer*. 2007;110:173-8.
19. Cren PY, Lebellec L, Ryckewaert T, Penel N. Anti-angiogenic agents in management of sarcoma patients: overview of published trials. *Front. Oncol*. 2020(10):594445.

The World Journal of Medical Education & Research (WJMER) is the online publication of the Doctors Academy Group of Educational Establishments. It aims to promote academia and research amongst all members of the multi-disciplinary healthcare team including doctors, dentists, scientists, and students of these specialties from all parts of the world. The journal intends to encourage the healthy transfer of knowledge, opinions and expertise between those who have the benefit of cutting-edge technology and those who need to innovate within their resource constraints. It is our hope that this interaction will help develop medical knowledge & enhance the possibility of providing optimal clinical care in different settings all over the world.



WJMER

World Journal of Medical Education and Research

An Official Publication of the Education and Research Division of Doctors Academy

